



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Substituting from the condition $nk^{2(n-1)} = 1$ we find

$$\frac{d^2T}{dk^2} = -\frac{d(n-1)}{k} \left(\frac{n-1}{a \log n} \right)^{1/2}.$$

Since this result is negative ($n > 1$) a true maximum obtains.

Case (b). $1 > n > 0$.

Then $\overline{FP} < \overline{FQ}$. Proceeding as before we get

$$T'' = - \int_{k^nd}^{kd} \left(2a \log \frac{d}{x} \right)^{-1/2} dx,$$

where T'' symbolizes the interval of time consumed in going from Q to P . Consequently $nk^{2(n-1)} = 1$ and

$$\frac{d^2T''}{dk^2} = -\frac{d(1-n)}{k} \left(\frac{1-n}{-a \log n} \right)^{1/2}.$$

Since $n - 1$ is now negative, the second derivative is also negative, and hence the condition for a maximum is again fulfilled.

Remark: In making the final reductions in case (b) attention has to be paid to the fact that $\log n$ is negative as well as $n - 1$.

Also solved by A. H. WILSON, H. N. CARLETON, H. POLISH, J. A. CAPARO, ELIJAH SWIFT, PAUL CAPRON, and the PROPOSER.

NUMBER THEORY.

209. (March, 1914.) Proposed by R. D. CARMICHAEL, University of Illinois.

Prove that the difference of the sixth powers of two integers cannot be the square of an integer.

SOLUTION BY ELIJAH SWIFT, University of Vermont.

In Carmichael's Diophantine Analysis, pp. 70, 71, the impossibility of the equation $x^3 + y^3 = 2^nz^3$ is proved. (The proof is valid whether y is positive or negative.)

We are to prove the impossibility of the equation $a^6 - b^6 = c^2$. We know that if $x^2 + y^2 = z^2$, then $x = 2mn$, $y = m^2 - n^2$, $z = m^2 + n^2$. (Loc. cit., p. 10.) We assume, of course, that a, b, c are all prime to each other; so are x, y, z also. Two cases present themselves:

$$(I) \quad a^3 = m^2 + n^2, \quad b^3 = m^2 - n^2, \quad c = 2mn.$$

$$(II) \quad a^3 = m^2 + n^2, \quad b^3 = 2mn, \quad c = m^2 - n^2.$$

Case I. Of the two integers, m and n , one is odd and the other even. Also these numbers are prime to each other. Consequently $m + n$ and $m - n$ are prime to each other, and since their product is a cube, each of them must be a cube also. If we set them equal to α^3 and β^3 respectively

$$m + n = \alpha^3, \quad m - n = \beta^3, \quad a^3 = m^2 + n^2 = \frac{\alpha^6 + \beta^6}{2}, \quad \text{or} \quad \alpha^6 + \beta^6 = 2a^3,$$

the impossibility of which was stated in the first paragraph.

Case II. Assume m even; the proof will apply equally well to the other case, n even. Since $2mn$ is a cube, $2m$ and n must each of them be a cube. Setting them equal to $8\alpha^3$ and β^3 , we have

$$m = 4\alpha^3, \quad n = \beta^3, \quad a^3 = 16\alpha^3 + \beta^6 \quad \text{or} \quad 2^4\alpha^6 = a^3 - \beta^6,$$

an equation which was proved to be impossible. Hence the equation $a^6 - b^6 = c^2$ cannot hold.

219. (June, 1914.) Proposed by R. D. CARMICHAEL, University of Illinois.

Determine whether it is possible for a polygon to have the number of its diagonals equal to a perfect fourth power.

SOLUTION BY ELIJAH SWIFT, University of Vermont.

If the number of sides of a polygon is n , the number of its diagonals is $n(n-3)/2$, which by the conditions of the problem must be a perfect fourth power.